

EGU22-10172

<https://doi.org/10.5194/egusphere-egu22-10172>

EGU General Assembly 2022

© Author(s) 2022. This work is distributed under the Creative Commons Attribution 4.0 License.



Hydrocarbons investigations from near-surface sediments of the north and northeastern Barents Sea shelves

Olesya Vidischeva¹, Elena Poludetkina¹, Evgeniya Basova¹, Elizaveta Dralina¹, Aleksandr Bogdanov¹, Elena Bakay¹, Irina Man'ko¹, Grigorii Akhmanov¹, and **Adriano Mazzini**²

¹Faculty of Geology, Department of Petroleum Geology, Lomonosov Moscow State University, Moscow, Russian Federation (vid6877@yandex.ru)

²Centre for Earth Evolution and Dynamics (CEED), University of Oslo, Oslo, Norway (adriano.mazzini@geo.uio.no)

The Russian portion of the Barents Sea shelf is the largest offshore zone in Russia with high petroleum potential. Numerous offshore oil and gas fields have been discovered in the southern part of the Barents Sea, however little is known about the northern and northeastern sectors. These regions were investigated during the TTR-19 and TTR-20 expeditions with the aim to characterize the gas type and content in the near-surface sediments and to identify potential fluid migration areas.

Sites for seafloor coring were selected based on the acquired geophysical data, targeting seafloor morphologies of obvious interest (e.g. pockmarks, faulted zones or tunnel valleys) or subsurface acoustic anomalies observed on the seismic profiles. Lithological composition and gas extracted from the sampled sediments were analyzed using gas chromatography, pyrolysis, mass-spectrometry.

Results of hydrocarbon (HC) gas molecular studies showed some differences between the northern and northeastern parts of the Barents Sea. Northeastern Barents Sea shelf sediments are characterized by low concentrations of methane up to 28 ppm, and a small amount of C₂₊ compounds. Northern Barents Sea shelf sediments have methane concentrations up to 69.8 ppm and the presence of C₂H₆, C₂H₄, C₃H₈ and C₃H₆ and, in a few cores, also C₄H₁₀ and C₅H₁₂. The study of the organic matter (OM) of bottom sediments (upper 2 meters) also showed a difference in the composition of its soluble part. The OM concentrations in northern part are higher than those observed in the northeastern part, and are characterized by the presence of light HC and oily compounds, which may indicate migration processes taking place in sedimentary covers. Geophysical studies conducted in the northern part, show that the complex of dense subglacial sediments is only locally distributed. These deposits are instead ubiquitous in the northeastern part and serve as a lithological barrier preventing the migration of fluids to the surface. Mass-spectrometry studies allowed the identification of the contemporary OM biomarker outlook. Hopanes and steranes with highly characteristic distributions of structural and stereochemical isomers (e.g. like in sediments with mature organic matter) were confidently identified in a few stations. In recent sediments, with poor thermal alteration, such as those studied in this research, organic matter with higher maturity can most likely be attributed to migration of thermogenic HCs.

Overall the bottom sediments collected in the northern and northeastern parts of the Barents Sea showed low concentrations of OM and low amounts of methane from the headspace analyses. These observations may argue against focused active HC seepage in the study areas, nevertheless the molecular and isotopic composition indicates the presence of thermogenic gas. Therefore a fluid migration from deeper units can be inferred. We suggest that the distinct lithological variations and properties of Arctic bottom sediments are responsible for the different compositions (gases and OM) observed in the northern and northeastern parts and for the formation of background and anomalous concentrations of fluids in the near-surface sediments.