Acta Crystallographica Section E Structure Reports Online

ISSN 1600-5368

Elena V. Karpova,^a* Maxim A. Zakharov,^b Sergey I. Gutnikov^c and Roman S. Alekseyev^a

^aInorganic Chemistry Division, Chemistry Department, Moscow State University, 119992 Leninskie Gory 1-3, Moscow, Russia, ^bGeneral Chemistry Division, Chemistry Department, Moscow State University, 119992 Leninskie Gory 1-3, Moscow, Russia, and ^cHigh School of Materials Science, Moscow State University, 119992 Leninskie Gory, Moscow, Russia

Correspondence e-mail: karpova@inorg.chem.msu.ru

Key indicators

Single-crystal X-ray study T = 293 K Mean σ (C–C) = 0.003 Å R factor = 0.053 wR factor = 0.134 Data-to-parameter ratio = 15.2

For details of how these key indicators were automatically derived from the article, see http://journals.iucr.org/e. The asymmetric unit of the title compound, $2C_2H_8N^+ \cdot C_8H_4O_4^{2-}$, comprises two crystallographically independent dimethylammonium cations and two half-terephthalate anions. The latter are each disposed about an inversion centre. N-H···O hydrogen bonds link the ions into a threedimensional framework.

Bis(dimethylammonium) terephthalate

Received 6 October 2004 Accepted 8 November 2004 Online 30 November 2004

Comment

The title compound, (I), was isolated as a side-product in the synthesis of metal-organic framework materials. The asymmetric unit (Fig. 1) comprises two crystallographically independent dimethylammonium cations and halves of two terephthalate anions, the latter each disposed about an inversion centre. Cations and anions are connected by typical $N-H\cdots$ O hydrogen bonds into a three-dimensional framework (Table 1). The structure consists of two independent interpenetrating frameworks, each comprising two kinds of cations and two kinds of anions connected by hydrogen bonds



Experimental

The typical quantities of reagents were 0.2 mmol neodimium chloride or holmium nitrate and 0.3 mmol of terephthalic acid in 5 ml of DMSO in one vessel and 2 ml of dimethylamine in another vessel. For a typical synthesis of metal-organic framework materials, two DMFA solutions were prepared in glass vessels: a metal salt with terephthalic acid and the other containing only dimethylamine. The former glass vessel had only a small hole, whereas the latter was open. Such a technique is needed for the slow diffusion of amine into the reaction vessel for slow crystallization of metal-organic framework materials. Glass vessels of these solutions were placed into a desiccator held under vacuum. After a few days, colourless needles were formed. These crystals were mounted in a glass capillary under water-free conditions.

Crystal data $2C_2H_8N^+ \cdot C_8H_4O_4^{-2}$ $D_x = 1.198 \text{ Mg m}^{-3}$ $M_r = 256.30$ Mo $K\alpha$ radiation Monoclinic, $P2_1/n$ Cell parameters from 20 a = 9.642 (2) Åreflections b = 11.103 (2) Å $\theta = 10 - 14^{\circ}$ $\mu = 0.09~\mathrm{mm}^{-1}$ c = 13.272 (3) Å $\beta = 91.01(3)^{\circ}$ T = 293 (2) K $V = 1420.6 (5) \text{ Å}^3$ Prism, colourless $0.5 \times 0.3 \times 0.2 \text{ mm}$ Z = 4

© 2004 International Union of Crystallography Printed in Great Britain – all rights reserved

organic papers

Data collection

Enraf–Nonius CAD-4 diffractometer Non-profiled ω scans Absorption correction: ψ scan (North *et al.*, 1968) $T_{min} = 0.968, T_{max} = 0.982$ 2910 measured reflections 2790 independent reflections 1759 reflections with $I > 2\sigma(I)$

Refinement

Table 1

Refinement on F^2 $R[F^2 > 2\sigma(F^2)] = 0.053$ $wR(F^2) = 0.134$ S = 1.052790 reflections 184 parameters H atoms treated by a mixture of independent and constrained refinement
$$\begin{split} R_{\rm int} &= 0.050 \\ \theta_{\rm max} &= 26.0^{\circ} \\ h &= -11 \rightarrow 11 \\ k &= 0 \rightarrow 13 \\ l &= 0 \rightarrow 16 \\ 2 \text{ standard reflections} \\ \text{frequency: } 7200 \text{ min} \\ \text{intensity decay: } 1\% \end{split}$$

$$\begin{split} &w = 1/[\sigma^2(F_o^2) + (0.0626P)^2 \\ &+ 0.1137P] \\ &where P = (F_o^2 + 2F_c^2)/3 \\ (\Delta/\sigma)_{max} = 0.001 \\ \Delta\rho_{max} = 0.20 \text{ e } \text{\AA}^{-3} \\ \Delta\rho_{min} = -0.23 \text{ e } \text{\AA}^{-3} \\ &\text{Extinction correction: SHELXL97} \\ &\text{Extinction coefficient: } 0.083 (7) \end{split}$$

Hydrogen-bonding geometry (Å, °).				
$D - H \cdots A$	D-H	$H \cdots A$	$D \cdots A$	$D - H \cdots A$
N1-H5···O3 ⁱ	0.95 (2)	1.83 (2)	2.761 (3)	167 (2)
$N1 - H6 \cdots O2$	0.96 (3)	1.78 (3)	2.731 (3)	171 (2)
$N2-H1\cdots O1$	0.95 (3)	1.77 (3)	2.700 (3)	165 (2)
$N2-H2\cdots O4^{ii}$	1.02 (3)	1.68 (3)	2.693 (3)	179 (3)

Symmetry codes: (i) 1 + x, y, z; (ii) $\frac{1}{2} + x, \frac{1}{2} - y, z - \frac{1}{2}$.

N-bound H atoms, *i.e.* H1, H2, H5 and H6, were located in a difference map and refined isotropically. As the N1–H5 distance was too long under free refinement, it was restrained to 0.95 (2) Å. C-bound H atoms were included in the riding-model approximation, with C–H = 0.93 Å (for benzene) and C–H = 0.96 Å (for methyl), and $U_{\rm iso}(\rm H) = 1.2U_{eq}(\rm CH)$ and $U_{\rm iso}(\rm methyl H) = 1.5U_{eq}(\rm CH_3)$.

Data collection: *CAD-4 EXPRESS* (Enraf–Nonius, 1994); cell refinement: *CAD-4 EXPRESS*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *DIAMOND* (Crystal Impact, 2000); software used to prepare material for publication: *WinGX* (Farrugia, 1999).



Figure 1

A view of the association between ions in dimethylammonium terephthalate, showing the atomic numbering scheme. Molecules of only one framework are shown. The second framework is hidden for simplicity. Displacement ellipsoids are drawn at the 50% probability level. [Symmetry codes: (i) $x + \frac{1}{2}, \frac{1}{2} - y, z - \frac{1}{2}$; (ii) x + 1, y, z.]

References

- Crystal Impact (2000). *DIAMOND*. Crystal Impact GbR, Postfach 1251, D-53002 Bonn, Germany.
- Enraf-Nonius (1994). CAD-4 EXPRESS. Enraf-Nonius, Delft, The Netherlands.
- Farrugia, L. J. (1999). J. Appl. Cryst. 32, 837-838.
- Harms, K. & Wocadlo, S. (1995). XCAD4. University of Marburg, Germany. North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). Acta Cryst. A24, 351– 359.
- Sheldrick, G. M. (1997). SHELXS97 and SHELXL97. University of Göttingen, Germany.